

Injury Reduction Effectiveness of Assigning Running Shoes Based on Plantar Shape in Marine Corps Basic Training

Joseph J. Knapik,^{*†} ScD, Daniel W. Trone,[‡] PhD(c), David I. Swedler,[§] MPH, Adriana Villasenor,[‡] MPH, Steve H. Bullock,[†] DPT, Emily Schmied,[‡] MPH, Timothy Bockelman,^{||} BEd, Peggy Han,[‡] MPH, and Bruce H. Jones,[†] MPH, MD
 From the [†]US Army Public Health Command, Aberdeen Proving Ground, Maryland, [‡]Naval Health Research Center, San Diego, California, [§]Johns Hopkins University, Baltimore, Maryland, and ^{||}Marine Corps Recruit Depot, Eastern Recruiting Region, Parris Island, South Carolina

Background: Shoe manufacturers market motion control, stability, and cushioned shoes for plantar shapes defined as low, normal, and high, respectively. This assignment procedure is presumed to reduce injuries by compensating for differences in running mechanics.

Hypothesis: Assigning running shoes based on plantar shape will not reduce injury risk in Marine Corps basic training.

Study Design: Randomized controlled clinical trial; Level of evidence, 1.

Methods: After foot examinations, Marine Corps recruits in an experimental group (E: 408 men, 314 women) were provided motion control, stability, or cushioned shoes for plantar shapes indicative of low, medium, or high arches, respectively. A control group (C: 432 men, 257 women) received a stability shoe regardless of plantar shape. Injuries during the 12 weeks of training were determined from outpatient visits obtained from the Defense Medical Surveillance System. Other known injury risk factors (eg, fitness, smoking, prior physical activity) were obtained from a questionnaire, existing databases, or the training units.

Results: Cox regression indicated little difference in injury risk between the E and C groups among men (hazard ratio [E/C] = 1.01; 95% confidence interval, 0.82-1.24) or women (hazard ratio [E/C] = 0.88; 95% confidence interval, 0.70-1.10).

Conclusion: This prospective study demonstrated that assigning shoes based on the shape of the plantar foot surface had little influence on injuries even after considering other injury risk factors.

Keywords: physical fitness; physical activity; tobacco use; prior injury; demographics; menstrual dysfunction

Injuries are a significant problem in the military and have a major effect on operational readiness because of medical costs, lost training time, and attrition associated with these injuries.⁴⁻⁶ Historically, injuries have been shown to be the leading causes of disability, hospitalization, and

*Address correspondence to Joseph J. Knapik, ScD, Directorate of Epidemiology and Disease Surveillance, US Army Public Health Command, Aberdeen Proving Ground, MD 21010 (e-mail: joseph.knapik@us.army.mil).

The authors declared that they had no conflicts of interest in their authorship and publication of this contribution. The use of trademarked name does not imply endorsement by any military service but is intended only to assist in identification of the characteristic of a specific product. All materials constitute the personal statements of the authors and are not intended to constitute an endorsement by any other federal government entity.

The American Journal of Sports Medicine, Vol. XX, No. X
 DOI: 10.1177/0363546510369548
 © 2010 The Author(s)

outpatient visits in the military services.^{5,8,28,29} In 2008, injuries and musculoskeletal conditions among members of active duty Armed Forces were responsible for 17 218 hospital admissions and over 2.7 million ambulatory visits.^{19,20} Injury-related conditions account for limited duty rates of 40 to 120 days per 100 soldiers per month.³² The occurrence of a training-related injury is associated with poor long-term military outcomes.^{12,31} Previous studies have found varying injury incidence, but generally, about 25% of men and 45% of women will experience one or more injuries during the course of Marine Corps basic training.^{1,2,7,24,27} The majority of overuse injuries involve the lower extremities and appear to come from physical fitness training activities, specifically running.³⁰

When recruits first arrive for Marine Corps basic training, they are issued a new pair of running shoes. At the time the current study was conducted, these shoes were assigned based on the foot surface area contacting the floor (ie, the shape of the plantar surface) while standing. The

plantar shape during static weightbearing was presumed to reflect foot arch height. Recruits judged to have high, low, and normal arches were assigned cushioned shoes, motion control shoes, or stability shoes, respectively. Shoe manufacturers market these 3 types of running shoes, and they are presumably designed to reduce injuries by compensating for differences in running mechanics. Cushioned shoes contain softer midsole material like ethyl vinyl acetate (EVA) that is thought to allow for greater shock absorption and permit more pronation for high-arched individuals who are assumed to impact the ground with high force and to underpronate. Motion control shoes are reputed to limit the excessive pronation experienced by lower arched individuals by using harder midsole materials (polyurethane) and other features that are specific to particular models. Stability shoes designed for normal-arched individuals are midway between motion control and cushioned type shoes, containing some motion control features and cushioning characteristics.²³

We previously reported on studies conducted in US Army Basic Combat Training¹⁶ and Air Force Basic Military Training¹⁰ showing that assigning running shoes on the basis of plantar shape did not reduce the risk of training-related injuries. The major purpose of the present study was to determine whether injury risk could be reduced in Marine Corps basic training (MCBT) by assigning running shoes based on the static weightbearing plantar foot shape.

MATERIALS AND METHODS

Participants

Participants were volunteers from among male basic trainees at the Marine Corps Recruit Depot (MCRD), San Diego, California, and female basic trainees at the MCRD, Parris Island, South Carolina. Potential volunteers were briefed on the purposes and risks of the study, and those wishing to participate signed an informed consent statement. The research protocols were approved by the Institutional Review Board of the Naval Health Research Center, San Diego, California.

Procedures

Immediately after informed consent was obtained, volunteers were administered a questionnaire that asked about tobacco use, physical activity, injury history, and (for women) menstrual history. Recruits then removed their shoes and socks and mounted the acrylic platform of the light box device.^{10,16} The device contained a mirror that reflected the underside of the trainee's foot. This provided a view of the footprint from above, showing how much of the foot was in contact with the acrylic surface. The participants were instructed to stand with equal weight on each foot, with their feet comfortably apart. The plantar surface area (footprint) was examined by trained evaluators who rated the plantar surface as either high arched, normal arched, or low arched, based on templates²³: more area

in contact with the acrylic surface in the middle third of the plantar surface indicated a low plantar shape, and less area in contact, a high plantar shape. In a subsample of cases ($n = 66$), 2 evaluators made independently recorded determinations of the plantar shape for the purposes of calculating between-rater reliability.

Following all foot measurements, participating recruits were randomized into 1 of 2 groups using an assignment order that was randomly generated by a statistical software program. Recruits assigned to the control (C) group received a stability shoe, the New Balance 767ST (Boston, Massachusetts), regardless of plantar shape. Recruits assigned to the experimental (E) group received a shoe based on the determined shape of the plantar surface of their foot. If the E-group recruit had a low arch, a motion control shoe, New Balance 587NV, was assigned. If the E-group recruit had a high arch, a cushion shoe, New Balance 881WG, was assigned. If the E-group recruit had a normal arch, a stability shoe was assigned, a New Balance 767ST. All participating recruits were asked their shoe size, and a shoe of this size was initially provided. If the shoe did not fit, different shoe sizes were tried until a proper fit was achieved.

Physical Characteristics, Physical Fitness, and Demographics

Additional data were obtained from an existing administrative data source routinely collected and maintained by the recruit training staff. These data included weight, height, and physical fitness test scores measured before the first day of recruit training. The recruits' weight and height were measured in socks, T-shirts, and shorts using a standard, calibrated mechanical physician's beam scale with a stadiometer.

The fitness test for the men consisted of 3 events: pull-ups, abdominal crunches, and a 1.5-mile run, conducted in that order. The fitness test for the women involved the same abdominal crunch and 1.5-mile run events that were performed by the men. However, instead of the pull-ups, women performed a flexed arm hang. Test events were administered by drill instructors using well-standardized procedures.

The Armed Forces Health Surveillance Center (AFHSC)²⁵ provided demographic data for the recruits from the Defense Manpower Data Center (DMDC). Recruit information obtained from the DMDC included date of birth, component (active Marine Corps or reserve Marine Corps), educational level, marital status, and race.

Marine Corps Recruit Training

United States Marine Corps recruit training consisted of 12 weeks of standardized military instruction for both male and female recruits. Men and women were trained by drill instructors of their own sex. Because of logistical and geographical reasons alone, the training schedules at the 2 training locations (ie, Parris Island and San Diego)

vary only in the training day in which events occur. The 12 weeks included 5 to 7 processing and forming days, and 70 training days, with no formal training conducted on Sundays. Running shoes were worn during processing and forming. There were about 40 miles of running during physical training in the 12 weeks.

Formal recruit training was divided into 3 phases of about 23 to 24 training days. Phase 1 used a progressive physical training program, which included general physical conditioning (14 sessions), pugil stick training (2 sessions), water survival skills training (4 days), and 5-km and 8-km conditioning marches (1 each). General physical conditioning exercises included running, calisthenics, obstacle courses, and circuit courses. There were 4 days when no physical training was scheduled. Running shoes and Marine Corps combat boots were alternated on the first 16 training days, but after that, running shoes were worn for 12 to 15 hours per week of physical training. Phase 1 also included classroom instruction on Marine Corps history, core values, leadership, ethics, first aid training, health and hygiene, personal appearance, and uniform instruction.

Phase 2 emphasized marksmanship fundamentals using the M-16A2 rifle and a final qualifying test. Physical activity included general physical conditioning (7 sessions), platoon drill, and 10-km and 12-km marches (1 each). Running shoes were worn for 8 to 10 hours per week of physical training. There were 9 days when no physical training was scheduled.

Phase 3 focused on a field training exercise (the Crucible), which took place over 54 hours, during training days 63 to 65. The first day of the exercise began with a 10-km march and ended with an 8-km night march. The last day finished with a 15-km march. Other physical training consisted of Basic Warrior Training (BWT, 4 sessions), general physical conditioning (4 sessions), platoon drills, and a motivation run the day before graduation. The BWT course was 2 one quarter miles with obstacles every quarter mile; one session was performed in running shoes. Running shoes were worn 8 to 10 hours per week in phase 3, and there were 6 days when no physical training was scheduled. The final few training days involved continued drill and ceremony, practice for graduation, and the graduation ceremony.

Attrition From Training

Some participants did not complete the entire 12-week basic training cycle, but their data were included for the time they remained in training, as described below. Reasons for attrition included discharge from the Marine Corps or reassignment to a new company (recycle). Discharges and recycles were obtained from a local data system maintained at MCRD San Diego or MCRD Parris Island.

Injury Outcome Measures

Injury outcome data were obtained from the AFHSC. The AFHSC systematically obtains data on ambulatory (outpatient) encounters that occur within military treatment facilities (MTFs) or outside MTFs but are paid for by the

Department of Defense. The AFHSC provided visit dates and International Classification of Diseases, Version 9, Clinical Modification (ICD-9-CM) codes for all outpatient medical visits within the recruit training time frame for each recruit. The first 4 diagnoses for each visit were considered, although a single visit usually included only one diagnosis. Five injury indices were calculated: the Installation Injury Index (III), Modified Installation Injury Index (MIII), Training-Related Injury Index (TRII), Overuse Injury Index (OII), and Comprehensive Injury Index (CII). These indices include specific ICD-9 codes, as described previously.¹³ The III has been used to compare overall injury rates (acute and overuse) among military posts and is reported on a monthly basis at the Armed Forces Health Surveillance Center Web site (<http://afhsc.army.mil>), where the ICD-9-CM codes are also provided. The MIII is similar to the III but captures a greater number of injuries than the III, including more overuse-type injuries. The TRII is limited to lower extremity overuse injuries and has been used to compare injury rates among Army basic training posts. The OII captures the subset of musculoskeletal injuries presumably resulting from cumulative microtrauma (overuse injuries) such as stress fractures, stress reactions, tendinitis, bursitis, fasciitis, arthralgia, neuropathy, radiculopathy, shin splints, synovitis, and musculoskeletal pain (not otherwise specified). The CII captures all ICD-9 codes related to injuries including overuse and traumatic.

Data Analysis

Data were compiled and analyzed using SPSS version 16.0.1 (Chicago, Illinois). Age was calculated from the date of birth in the DMDC data to the date of the informed consent briefing. Body mass index (BMI) was calculated as weight/height² (kg/m²).¹¹

The E and C groups were compared on attrition, age, physical characteristics, physical fitness, demographic characteristics, and the questionnaire variables. For continuous measures, group comparisons were performed using an independent sample *t* test; for discrete, nominal, and ordinal variables, comparisons were made using the χ^2 statistic. Between-rater reliability of plantar foot shape determination was made with the κ coefficient.

Person-time injury incidence rates (injured subjects/1000 person-days) for all injury indices were calculated as $([\text{recruits with } \geq 1 \text{ injury by index}]/[\text{total recruit time in basic training}]) \times 1000$. Comparisons between the E and C groups were made using a χ^2 for person-time.⁹

Cox regression (survival analysis) was used to examine group differences in time to first injury, examining the OII, TRII, and CII. The number of days in training was the time scale. Univariate Cox regression involved group (E or C) as the only independent variable. The C group was defined as the reference group (hazard ratio = 1.00), and simple contrasts were made between the C and E groups. Multivariate Cox regression included any variable that differed between the groups in the χ^2 or *t* test analyses ($P < .10$). For each Cox regression analysis, once a recruit had an




Figure 1. Flow of volunteers through study.

injury, his or her contribution to time in MCBT was terminated. Those who withdrew from MCBT had their times censored (ie, end of time at risk) at the day they left training.

RESULTS

Participants, Attrition, and Reliability

In total, 917 men and 694 women volunteered for the study. Not considered in the analyses were recruits who left the testing area without receiving a group assignment (due to externally imposed time constraints), those not entering basic training, those who did not have a graduation date in administrative records, and/or those whose data were not obtained from the AFHSC. The final cohort consisted of 840 men and 571 women. The initial volunteers and loss of participants by group is shown in Figure 1.

Among the men, 15.3% in the C group and 11.5% in the E group withdrew from training ($P = .11$). Among the women, 12.1% of the C group and 11.5% of the E group withdrew from training ($P = .83$). The κ coefficient comparing 2 raters on the plantar surface evaluations was 0.91 for the right foot and 0.91 for the left foot ($n = 66$), suggesting strong agreement and high comparative quality beyond chance alone.²⁶

Comparisons of C and E Groups

Sample sizes shown in the tables below can be different depending on the completeness of the data. This occurred because not all participants had complete measurements on all variables: a recruit may not have provided a response on the questionnaire or the AFHSC or the training unit may not have had some specific information.

Appendix 1 (available in the online version of this article at <http://ajs.sagepub.com/supplemental/>) compares group differences in age, physical characteristics, fitness scores, and the questionnaire items with continuous numeric responses. Group differences in age and the physical characteristics were small for both men and women. With regard to physical fitness, C-group men performed more pull-ups and crunches compared with the E-group men, but fitness differences were small between the C- and E-group women. None of the questionnaire items in Appendix 1 differed significantly between the C- and E-group men or women.

Appendix 2 (available in the online version of this article at <http://ajs.sagepub.com/supplemental/>) shows group comparisons on the demographic variables and the questionnaire items with ordinal, nominal, and discrete responses. The distribution of participants was similar for the C and E groups on the demographic variables for both men and women. On the questionnaire items, the E-group men had more recruits reporting that they had

TABLE 1
Comparison of Injury Incidence Rates Between Experimental (E) and Control (C) Groups^a

Injury Index	Men				Women			
	Injury Incidence Rate (injuries/1000 person-days)		Rate Ratio-C/E (95% CI)	P Value ^b	Injury Incidence Rate (injuries/1000 person-days)		Rate Ratio-C/E (95% CI)	P Value ^b
	C (n = 432)	E (n = 408)			C (n = 257)	E (n = 314)		
Installation	5.14	5.40	0.95 (0.77-1.19)	.70	5.43	4.57	1.19 (0.91-1.55)	.24
Modified Installation	5.24	5.40	0.97 (0.78-1.21)	.79	5.90	4.59	1.20 (0.93-1.56)	.19
Overuse	4.14	4.06	1.02 (0.79-1.31)	.89	3.29	2.80	1.18 (0.83-1.66)	.41
Training-Related	3.56	3.63	0.98 (0.75-1.28)	.88	1.49	2.03	1.18 (0.79-1.77)	.48
Comprehensive	5.72	5.76	0.99 (0.80-1.22)	.95	6.00	4.96	1.21 (0.94-1.57)	.16

^aCI, confidence interval.
^b χ^2 statistic for person-time.⁹

TABLE 2
Injury Risk in the Control (C) and Experimental (E) Groups (Cox Regression)^a

Injury Index	Group	Men (Univariate Analysis)		Women (Univariate Analysis)		Men (Multivariate Analysis) ^b	
		Hazard Ratio-C/E (95% CI)	P Value	Hazard Ratio-C/E (95% CI)	P Value	Hazard Ratio-C/E (95% CI)	P Value
Overuse	Control	1.00	—	1.00	—	1.00	—
	Experimental	0.99 (0.76-1.27)	.96	0.84 (0.59-1.19)	.32	1.03 (0.80-1.34)	.81
Training-Related	Control	1.00	—	1.00	—	1.00	—
	Experimental	1.04 (0.80-1.36)	.77	0.84 (0.56-1.26)	.39	1.07 (0.81-1.41)	.65
Comprehensive	Control	1.00	—	1.00	—	1.00	—
	Experimental	1.01 (0.82-1.24)	.94	0.88 (0.70-1.10)	.15	1.06 (0.85-1.32)	.62

^aCI, confidence interval.
^bIncludes pull-up performance, crunch performance, cigarette smoking, and prior lower limb injury.

smoked 100 cigarettes in their lifetime and more recruits reporting a prior lower limb injury. The E- and C-group women were similarly distributed on the questionnaire responses.

Injuries, Injury Rates, and Injury Risk

The 10 most common ICD-9-CM codes for the men were the following (in order of high to lower frequency): 726.69 (enthesopathy of knee), 848.8 (other sprains/strains), 845.00 (sprains/strains of ankle and foot), 844.9 (sprains/strains of knee and leg), 717.9 (internal derangement of knee), 919 (superficial injury of other, multiple, and unspecified sites), 847.9 (sprains and sprains of unspecified back sites), 719.46 (pain in lower leg), 840.9 (sprain/strain of shoulder and upper arm), and 843.9 (sprain/strain of hip and thigh). The 10 most common ICD-9-CM codes for the women were the following (in order of high to lower frequency): 729.5 (pain in limb), 919 (superficial injury of other, multiple, and unspecified sites), 719.46 (pain in lower leg), 719.45 (pain in pelvic region and thigh), 845.00 (sprains/strains of ankle and foot), 719.47 (pain in

ankle and foot), 719.41 (pain in shoulder region), 848.9 (sprains/strains, unspecified site), 919 (superficial injury of other, multiple, and unspecified sites), and 728.71 (plantar fascial fibromatosis).

Table 1 shows the person-time injury incidence rates for the various injury indices and compares the rates in the C and E groups. The incidence rates for the C and E groups were similar for both men and women for all of the indices.

Table 2 shows the results of the Cox regressions. Univariate analyses showed that for both men and women, there was little difference in OII, TRII, or CII risk between the groups. Because the 2 groups of men differed on pull-ups, crunches, smoking, and prior lower limb injury, a multivariate Cox regression was performed including these measures as covariates with the group factor. The hazard ratios for the men changed little when these covariates were included in a multivariate Cox regression.

Table 3 shows Cox regressions comparing CII risk between groups with high and low plantar shapes who wore different shoe types. Among the men and women with either low or high plantar shapes, injury risk was similar regardless of shoe type.

TABLE 3

Comparison of Recruits With Low and High Plantar Shapes Wearing Different Shoe Types (Comprehensive Injury Index)^a

Plantar Shape	Shoe Comparison	Men			Women		
		n	Hazard Ratio-E/C (95% CI)	P Value	n	Hazard Ratio-E/C (95% CI)	P Value
Low	Motion control/stability	62	0.91 (0.40-2.07)	.82	57	0.74 (0.31-1.76)	.49
High	Cushion/stability	79	1.05 (0.53-2.10)	.89	105	1.11 (0.62-2.00)	.72

^aE, experimental group; C, control group; CI, confidence interval.

TABLE 4

Running Shoes Used in the Marine Corps, Air Force, and Army Physical Training Footwear Studies

Service	Experimental Group Shoes			
	Motion Control Shoe	Stability Shoe	Cushion Shoe	Control Group Stability Shoe
Marine Corps	New Balance 587	New Balance 767	New Balance 881	New Balance 767
Air Force	New Balance 587	New Balance 498	New Balance 755	New Balance 498
Army	Asics Gel Foundation 7	Asics Gel 1120	Asics Gel Cumulus	New Balance 767
	Brooks Addiction 7	Asics Gel 2120	Brooks Radius 6	
	Saucony Grid Stabil 6	Brooks Adrenaline GTS6	Nike Air Pegasus	
	New Balance 857 ^a	Brooks Adrenaline GTS7	Saucony Grid Trigon 4	
		Nike Structure Triax	New Balance 644	
		Nike Air Max Moto ^a	New Balance 755	
		Saucony Grid Omni 5		
	New Balance 717			
	New Balance 767			

^aFor 2 shoes, the Army classification differed from those of the Runner's World and the manufacturer. One shoe was the New Balance 857, which the Army classification listed as a motion control shoe but Runner's World and the manufacturer listed as a stability shoe; the other was the Nike Air Max Moto, listed in the Army classification as a stability shoe but by Runner's World and the manufacturer as a cushioned shoe.

DISCUSSION

The present study demonstrated that assigning running shoes on the basis of the shape of plantar foot surface did not reduce injury risk in MCBT. Men and women who wore the shoe assigned based on their plantar shape had similar injury risk when compared with those who received a standard stability shoe regardless of foot type.

The results of the current study can be compared with the results of similar Army¹⁶ and Air Force¹⁰ basic training investigations, which also examined the effectiveness of assigning shoes based on plantar shape. These studies were designed to be complementary, but there were some important differences. Similarities among the 3 studies included (1) tracking participants in the same medical surveillance system, (2) calculating injury incidence rates in an identical manner, (3) an identical lifestyle questionnaire, and (4) the same randomized prospective design with the C group receiving a single stability shoe and an E group receiving a shoe based on plantar shape. Major differences among the Services' studies had to do with the brands and models of the shoes provided and the nature of the training environment. Table 4 shows the shoes used in the Marine Corps, Air Force, and Army

investigations. The C-group participants in the Marine Corps and Army studies received the same stability-type shoe, but C-group participants in the Air Force investigation received another type of stability shoe. The E-group participants in the Marine Corps and Air Force study received only 1 of 3 shoes, one for each foot type; E participants in the Army study could select from 19 different shoes, as long as the shoe they chose had been designated as appropriate for their plantar shape. There are also differences in the Marine Corps, Air Force, and Army basic training programs of instruction and length of training (12, 6, and 9 weeks, respectively).

Despite the differences in the Marine Corps, Air Force,¹⁰ and Army¹⁶ studies, the results generally concurred in showing that assigning running shoes based on plantar shape had little influence on injury risk during basic training. Hazard ratio comparisons for the 3 studies are shown in Table 5. To more fully examine injury risk when shoes were assigned based on plantar shape, meta-analysis was performed. The meta-analysis method was a general variance-based technique that employed univariate hazard ratios and confidence intervals from each study to produce a summary hazard ratio (SHR) and summary 95% confidence interval (S95% CI) that reflected the

TABLE 5
Meta-Analysis of Studies Examining if Injury Risk Can be Reduced by Assigning Shoe Based on Plantar Shape (Comprehensive Injury Index)^a

Service	Men		Women	
	Hazard Ratio-E/C (95% CI)	Summary Hazard Ratio-E/C (S95% CI)	Hazard Ratio-E/C (95% CI)	Summary Hazard Ratio-E/C (S95% CI)
Marine Corps	1.01 (0.82-1.24)	1.04 (0.94-1.14)	0.88 (0.70-1.10)	1.05 (0.95-1.18)
Air Force	1.09 (0.92-1.29)		1.23 (1.00-1.53)	
Army	1.02 (0.89-1.17)		1.06 (0.90-1.24)	

^aE, experimental group; C, control group; 95% CI, 95% confidence interval; S95% CI, summary 95% confidence interval.

TABLE 6
Meta-Analysis of Recruits With Low and High Plantar Shapes Wearing Different Shoe Types (Comprehensive Injury Index)^a

Service	Plantar Shape	Shoe Comparison	Men		Women	
			Hazard Ratio-E/C (95% CI)	Summary Hazard Ratio-E/C (S95% CI)	Hazard Ratio-E/C (95% CI)	Summary Hazard Ratio-E/C (S95% CI)
Marine Corps	Low	Motion control/stability	0.91 (0.40-2.07)	Low: 1.13 (0.85-1.51) High: 1.19 (0.93-1.52)	0.74 (0.31-1.76)	Low: 1.11 (0.75-1.65) High: 1.12 (0.85-1.48)
	High	Cushion/stability	1.05 (0.53-2.10)		1.11 (0.62-2.00)	
Air Force	Low	Motion control/stability	1.33 (0.80-2.21)		0.95 (0.47-1.93)	
	High	Cushion/stability	1.01 (0.66-1.55)		1.41 (0.77-2.58)	
Army	Low	Motion control/stability	1.08 (0.73-1.60)		1.48 (0.83-2.63)	
	High	Cushion/stability	1.36 (0.97-1.91)		1.04 (0.72-1.49)	

^aE, experimental group; C, control group; 95% CI, 95% confidence interval; S95% CI, summary 95% confidence interval.

combined results of all the studies.²² As shown in Table 5, the SHRs indicated that there was little difference in injury risk between the C and E groups for men or women. Said another way, there was little difference in injury risk if the subjects received a stability shoe (C group) or received a shoe based on plantar surface (E group) after combining the results of the Marine Corps, Air Force, and Army studies.

As noted earlier, motion control shoes are designed for low-arched individuals to presumably control for excessive pronation; cushioned shoes are designed for high-arched individuals to presumably provide cushioning to reduce ground-impact forces and to allow for more foot pronation.^{17,18,21,33} If injury risk could be reduced by assigning running shoes based on plantar shape, that reduced risk might be best seen by comparing E and C groups with low or high plantar shapes. This is because the low plantar-shape E group wore motion control shoes designed by shoe companies for their foot type, but the low plantar-shape C group wore stability shoes designed for another foot type. Likewise, the high plantar-shape E group wore cushioned shoes designed by shoe companies for their plantar shape, but high plantar-shape C group participants wore stability shoes designed for another foot type. Table 6 shows these comparisons in the Marine Corps, Army, and Air Force studies for men and women separately. Again, meta-analysis employing a general variance-based technique was used to combine the results of the 3 studies,

and the SHR and S95% CI indicated the combined results.²² Contrary to expectation, SHRs in Table 6 indicated that injury risk was modestly elevated in the E group after combining the Marine Corps, Air Force, and Army studies.

Despite the general concurrence among the Marine Corps, Air Force,¹⁰ and Army¹⁵ investigations, these studies are not in accord with a previous Army study¹⁴ that showed a postwide decrease in serious injuries at Fort Drum, New York, after initiation of a running shoe prescription program. Methodological differences between the Fort Drum project and the current Marine Corps study are similar to those previously outlined in the Air Force and Army investigations.^{10,15} The current Marine Corps study involved a prescription based only on plantar shape; the Fort Drum project involved a prescription based on an evaluation of foot arch height and foot flexibility. The current Marine Corps study involved a population of recruits in a situation where there was assurance that the correct shoe was given to the recruit and worn during training. The Fort Drum project involved soldiers who were given the shoe prescription, but there was little follow-up to determine whether they had actually purchased and/or worn the recommended shoe. In fact, a survey involving a convenience sample of 122 Fort Drum soldiers (of an average 9752 estimated to be on post) found that only 11% self-reported that they had followed the shoe prescription advice. The current Marine Corps study involved

a prospective shoe prescription involving 2 randomly assigned groups (C and E) training side by side in a standardized program with follow-up for any injury occurring during the period. The Fort Drum project involved a retrospective examination of medical visits to a physical therapy clinic before and after the shoe program was initiated. A number of temporal factors were potential confounders in the Fort Drum project, and these were discussed at length in the report on that study.¹⁴ The major potential bias was a change in the medical surveillance system used to track injuries, which was discovered after investigating the time point when injuries dramatically decreased. In summary, the advantages of the current Marine Corps study were that it (1) involved a randomized prospective design manipulating only one variable (running shoe prescription based on plantar shape), (2) provided considerably better knowledge about the shoes worn, and (3) involved a more controlled training environment. Men and women in the current Marine Corps study trained in separate locations (San Diego, California, and Parris Island, South Carolina); however, the male and female data were analyzed separately, ensuring that gender-specific C and E groups were in the same training environment.

In conclusion, this prospective study demonstrated that assigning running shoes based on the static weightbearing plantar foot surface shape had little influence on injury risk during MCBT, even after controlling for other injury risk factors. The findings are strengthened by the similar results in Air Force and Army basic training studies and the meta-analysis reported above. If the goal is injury prevention, it is not necessary to provide running shoes to Marine Corps recruits based on a visual inspection of the static weightbearing plantar shape. This assignment procedure was no more protective against injury than issuing a single stability shoe regardless of plantar shape. It is still recommended that recruits receive a new shoe on entry to recruit training because older shoes have previously been shown to be associated with increased injury risk.³

ACKNOWLEDGMENT

This research was supported by the Naval Health Research Center, Work Unit No. 60626. We express our appreciation to the following members of our research team from the Naval Health Research Center in San Diego, California: Carol Macera, Mitchell Rauh, and Richard Shaffer. Additional thanks are due to James Reading from MCRD, San Diego, California.

REFERENCES

- Almeida SA, Trone DW, Leone DM, Shaffer RA, Patheal SL, Long K. Gender differences in musculoskeletal injury rates: a function of symptoms reporting? *Med Sci Sports Exerc.* 1999;31:1807-1812.
- Almeida SA, Williams KM, Shaffer RA, Brodine SK. Epidemiological patterns of musculoskeletal injuries and physical training. *Med Sci Sports Exerc.* 1999;31:1176-1182.
- Gardner LI, Dziados JE, Jones BH, et al. Prevention of lower extremity stress fractures: a controlled trial of a shock absorbent insole. *Am J Public Health.* 1988;78:1563-1567.
- Jones BH, Amoroso PJ, Canham ML, Weyandt MB, Schmitt JB. Atlas of injuries in U.S. Armed Forces. *Mil Med.* 1999;164(Suppl):1-1-9-25.
- Jones BH, Canham-Chervak M, Canada S, Mitchener TA, Moore S. Medical surveillance of injuries in the U.S. military: descriptive epidemiology and recommendations for improvement. *Am J Prev Med.* 2010;38(Suppl 1):S42-S60.
- Jones BH, Hansen BC. *Injuries in the Military: A Hidden Epidemic.* Technical Report No. 29-HA-4844-97. Aberdeen Proving Ground, Maryland: US Army Center for Health Promotion and Preventive Medicine; 1996.
- Jones BH, Shaffer RA, Snedecor MR. Chapter 6. Injuries treated in outpatient clinics: surveys and research data. *Mil Med.* 1999;164(8 Suppl):1-89.
- Jones JH, Perrotta DM, Canham-Chervak ML, Nee MA, Brundage JF. Injuries in the military: a review and commentary focused on prevention. *Am J Prev Med.* 2000;18(3S):71-84.
- Kahn HA, Sempos CT. *Statistical Methods in Epidemiology.* New York: Oxford University Press; 1989.
- Knapik JJ, Brosch LC, Venuto M, et al. Effect on injuries of assigning shoes based on foot shape in Air Force Basic Training. *Am J Prev Med.* 2010;38:S197-S211.
- Knapik JJ, Burse RL, Vogel JA. Height, weight, percent body fat and indices of adiposity for young men and women entering the U.S. Army. *Aviat Space Environ Med.* 1983;54:223-231.
- Knapik JJ, Canham-Chervak M, Hauret K, Hoedebecke E, Laurin MJ, Cuthie J. Discharges during US Army Basic Combat Training: injury rates and risk factors. *Mil Med.* 2001;166:641-647.
- Knapik JJ, Darakjy S, Scott S, et al. *Evaluation of Two Army Fitness Programs: The TRADOC Standardized Physical Training Program for Basic Combat Training and the Fitness Assessment Program.* Technical Report No. 12-HF-5772B-04. Aberdeen Proving Ground, Maryland: US Army Center for Health Promotion and Preventive Medicine; 2004.
- Knapik JJ, Feltwell D, Canham-Chervak M, et al. *Evaluation of Injury Rates During Implementation of the Fort Drum Running Shoe Injury Prevention Program.* Technical Report No. 12-MA-6558-01. Aberdeen Proving Ground, Maryland: US Army Center for Health Promotion and Preventive Medicine; 2001.
- Knapik JJ, Swedler D, Grier T, et al. *Injury Reduction Effectiveness of Prescribing Running Shoes Based on Foot Shape in Basic Combat Training.* Technical Report No. 12-MA-05SB-08. Aberdeen Proving Ground, Maryland: US Army Center for Health Promotion and Preventive Medicine; 2008.
- Knapik JJ, Swedler D, Grier T, et al. Injury reduction effectiveness of prescribing running shoes based on plantar shape. *J Strength Cond Res.* 2009;23:685-697.
- McPoil TG. Footwear. *Phys Ther.* 1988;68:1857-1865.
- McPoil TG. Athletic footwear: design, performance and selection issues. *J Sci Med Sport.* 2000;3:260-267.
- Medical Surveillance Group. Ambulatory visits among members of active components, U.S. Armed Forces, 2008. *Medical Surveillance Monthly Report.* 2009;16(4):10-15.
- Medical Surveillance Group. Hospitalizations among members of active components, U.S. Armed Forces, 2008. *Medical Surveillance Monthly Report.* 2009;16(4):2-8.
- Nigg BM, Segesser B. Biomechanical and orthopedic concepts in sports shoe construction. *Med Sci Sports Exerc.* 1992;24:595-602.
- Petitti DB. *Meta-Analysis, Decision Analysis and Cost-Effectiveness Analysis.* New York: Oxford University Press; 2000.
- Pritchard AE. Running shoe design, selection and care: does it make a difference? *US Army Med Dep J.* 2001;Apr/May/Jun:43-51.
- Rauh MJ, Macera CA, Trone DW, Shaffer RA, Brodine SK. Epidemiology of stress fractures and lower extremity overuse injuries in female recruits. *Med Sci Sports Exerc.* 2006;38:1571-1577.
- Rubertone MV, Brundage JF. The Defense Medical Surveillance System and the Department of Defense Serum Repository: a glimpse of the future of public health surveillance. *Am J Public Health.* 2002;92:1900-1904.

26. Seigel DG, Podgor MJ, Remaley NA. Acceptable values of kappa for comparison of two groups. *Am J Epidemiol.* 1992;135:571-578.
27. Shaffer RA, Brodine SK, Ito SI, Le AT. Epidemiology of illness and injury among U.S. Navy and Marine Corps female training populations. *Mil Med.* 1999;164:17-21.
28. Smith GS, Dannenberg AL, Amoroso PJ. Hospitalizations due to injuries in the military: evaluation of current data and recommendations on their use for injury prevention. *Am J Prev Med.* 2000;18(Suppl 3):41-53.
29. Songer TJ, LaPorte RE. Disabilities due to injury in the military. *Am J Prev Med.* 2000;18(Suppl 3):33-40.
30. Trank TV, Ryman DH, Minagawa RY, Trone DW, Shaffer RA. Running mileage, movement mileage, and fitness in male US Navy recruits. *Med Sci Sports Exerc.* 2001;33:1033-1038.
31. Trone DW, Villasenor A, Macera CA. Negative first-term outcomes associated with lower extremity injury during recruit training among female Marine Corps graduates. *Mil Med.* 2007;172:83-89.
32. US Army Disability Agency. *Disability Cost Estimates.* Department of the Army Inspector General's Report. Washington, DC: US Army Physical Disability Agency; 1995.
33. Winter DA, Bishop PJ. Lower extremity injury: biomechanical factors associated with chronic injury to the lower extremity. *Sports Med.* 1992;14:149-156.